



**RVC Exotics Service**  
**Beaumont Sainsbury Animal Hospital**  
**Royal Veterinary College**  
Royal College Street  
London  
NW1 0TU  
T: 0207 387 8134

---

---

---

Continuous egg laying in pet birds can be a significant problem leading to life threatening conditions such as depletion of calcium, egg binding (eggs stuck in the reproductive tract), reproductive infections and behavioural problems such as aggression. The most comm0 1 .V(8)11(t)()12(n)14(P(o)5)20( 13( )5)14(d(p)14(l)231 0 0057 m4(l)23)9)14(i)9)11( )1 0 0-121(i)

Reviewed November 2023

Removing eggs can stimulate the bird to continue to lay, as they  
m in place to allow her to brood them or replacing them with dummy  
eggs can help interrupt laying. Once the bird loses interest the eggs can be removed.

Medical or surgical therapy can be used to reduce egg laying in combination with these other interventions. Medical therapy consists of drugs designed to reduce the hormone output from the bird's ovary. This has traditionally been either as by injection (lasts 2-3 weeks) or an implant (can last for several months). Surgical therapy may be recommended to remove the oviduct and prevent egg laying, however there are risks and possible complications associated with this, so it is only considered after all other options have failed.

Seed is deficient in vitamins and minerals such as calcium, vitamin D and vitamin A. Birds should be gradually changed over to a complete pelleted diet, such as Harrison's or Nutriberries. This should comprise approximately 60-80% of the diet and 20-40% of the diet should be composed of high quality commercial seed. The diet should be supplemented with a complete vitamin and mineral supplement. This must be provided with the diet.

For calcium and vitamin D levels. This must be provided with the diet.

---

of the bird's diet. This must be provided with the diet.